
Socio-emotional outcomes of children in out of home care

Findings from the Pathways of Care Longitudinal Study (POCLS)

Professor Raghu Lingam and Dr Nan Hu on behalf of the Population Child Health Research Group



The Population Child Health Research Group



A multidisciplinary health systems research team working across the health services of NSW and University of New South Wales (UNSW). Our vision is to:

- Work with children and young people to discover the factors affecting their health, wellbeing, and access to the care they need.
- II. Conduct translational research that focuses on sustainably tailoring health systems to the needs of children and young people.
- III. Bridge the divide between research and service delivery by supporting staff to upskill in research and implementation of scalable interventions.

Children and Young People in Out-of-Home Care (OOHC)

- Children who are in the care of the local authority or who are provided with accommodation by the local authority social services, mostly as a result of abuse and/or neglect.
- In 2019, there were 45,000 children and young people (0-17 years) in OOHC in Australia. This number has been growing.
- These children have been identified as at high risk for many health difficulties including mental health difficulties.
- These mental health difficulties have been associated with exposure to maltreatment, age at entry into care, placement instability, caregiver characteristics, and relationships with caregivers.
- We need more evidence on which of these factors has the most influence. This will
 enable practitioners to better target their limited resources towards the right factors to
 improve outcomes for children and young people in OOHC.



The Pathways of Care Longitudinal Study (POCLS)

- The first large-scale prospective longitudinal study of children and young people in OOHC in Australia (hosted by the NSW Department of Communities & Justice).
- Links data from multiple agencies on child protection backgrounds, OOHC placements, health, education, and offending with first-hand accounts from children, caregivers, caseworkers, and teachers.
- Enables analysis of children in OOHC in general, and of priority populations including Aboriginal children and young people, children and young people from culturally and linguistically diverse backgrounds, and children and young people with high needs and/or disabilities.
- Provides a strong evidence base to inform policy, practice, and professional development to improve decision making and support provided to children and young people who cannot live safely at home.



Study aim

 To identify children in OOHC who are at high risk of socioemotional difficulties, both in early years after entry into OOHC and persistently over time, based on their demographic, precare, placement, and caregiver-related factors, using the POCLS data.




Factors examined in this study

- (i) child demographic factors, including age at interview, sex, Aboriginal status, and cultural background;
- (ii) pre-care maltreatment (i.e., maltreatment before first entry into OOHC), including child's age at the first entry into OOHC, number of unsubstantiated and substantiated reports of risk of significant harm (ROSH), respectively, and the predominant type of maltreatment;
- (iii) placement-related factors, including number of placement changes and duration in OOHC prior to the first interview, and placement type reported at the first interview;
- (iv) carer-related factors, including carer (and their spouse if recorded) socio-demographics (i.e., education level; financial situation), medical conditions, psychological distress measured using the Kessler Psychological Distress Scale (K10) (Kessler et al., 2002), and their satisfaction with support received from OOHC organisations.



Assessment of socio-emotional difficulties

- Children's socio-emotional outcomes were assessed at the first interview for each child (predominantly occurred within three years of first entry into OOHC)
 - the Brief Infant-Toddler Socio-emotional Assessment (BITSEA) for children aged 1-2 years
 - the Child Behaviour Check List (CBCL) for children aged 3-17 years





Children in OOHC at high risk of socio-emotional difficulties

- The overall proportion of children identified as having socio-emotional difficulties was 21.7% (95% CI, 19.6, 24.0).
- Two "high-risk" groups for socio-emotional difficulties

36% of children had socio-emotional difficulties if they

- are 3-5 years old at the POCLS interview, and
- live with a carer with high psychological stress, and
- 4+ placement changes


39% of children had socio-emotional difficulties if they

- are 6-17 years old at the POCLS interview, and
- live with a carer with high psychological stress, and
- experienced any substantiated allegation of child abuse, and
- were subject to emotional abuse alone or multiple types of abuse



Developmental trajectories of socioemotional difficulties of children in OOHC

- 345 children who participated in POCLS Waves 1 to 4 and who were aged 3 years or older at Wave 1 (after removing 2 children with a missing CBCL score).
- In the CBCL-clinical group, there were more children who
 - experienced 3 or more placements (58-75% increased risk), and
 - had a carer with a moderate to a high level of psychological distress (2.3 times increased risk), compared to children in the 'Persistently low difficulties' and the 'Normal' groups combined.





Study limitations

Cross-sectional

Sample size

Under-representation

- Study sample reflects younger children on long-term OOHC and protection orders. It under-represents adolescents aged 13-17 years who lived at home for a longer period before coming into OOHC.
- Thus, socio-emotional difficulties of the study sample may be lower than that of population who enter into OOHC and those who received final orders.



Overall, our study showed that 1 in 5 children aged 12 months to 17 years had socio-emotional difficulties predominantly within three years of entering OOHC.



This study identified five factors that identify children in OOHC who are at *high* risk of socio-emotional difficulties:

- age at assessment,
- living with a carer with high psychological distress,
- pre-care exposure to substantiated allegations,
- subject to multiple types of maltreatment, and
- 4+ placement changes.



This study should not distract child protection services from the need to provide health screening, and sufficient resources and support to all children in OOHC.

The current study does not take away from the need for individual level screening but rather adds a more nuanced risk profile of factors such as pre-care maltreatment and placement experiences that can inform early intervention and support for children in care.



Carer distress is an important indicator for the socio-emotional difficulties of the child they care that should be responded to with urgent support and interventions.

This finding highlights the importance of using a holistic approach to provide mental health support to both carers and children in need.




How can we help?

Need to think about models of care that will change practice

"stop describing start changing"



Supporting Looked After Children and Care Leavers in Decreasing Drugs and Alcohol (SOLID)



- A NIHR PHR research project that explores ways to decrease drug and alcohol use in children in OOHC and in care leavers aged 12-20 years in northeast England.
- References of articles by Alderson & Lingam et al.
- Alderson H, Kaner E, McColl E, Howel D, Fouweather T, et al. (2020) A pilot feasibility randomised controlled trial of two behaviour change interventions compared to usual care to reduce substance misuse in looked after children and care leavers aged 12-20 years: The SOLID study. PLOS ONE 15(9): e0238286. https://doi.org/10.1371/journal.pone.0238286
- Alderson, H, Brown, R, Smart, D, Lingam, R, Dovey-Pearce, G. 'You've come to children that are in care and given us the opportunity to get our voices heard': The journey of looked after children and researchers in developing a Patient and Public Involvement group. Health Expect. 2019; 22: 657- 665. https://doi.org/10.1111/hex.12904



Next steps

Priority population research

Aboriginal led research

OOHC research intervention based



Thank you!

